

4-1-2010

# An Approaching Epidemic: Exploring Diabetes Awareness & Care in Mombasa, Kenya

Benjamin Marhan Dropkin  
*SIT Study Abroad*, bdropkin@hamilton.edu

Follow this and additional works at: [http://digitalcollections.sit.edu/isp\\_collection](http://digitalcollections.sit.edu/isp_collection)

 Part of the [Public Health Education and Promotion Commons](#)

---

## Recommended Citation

Dropkin, Benjamin Marhan, "An Approaching Epidemic: Exploring Diabetes Awareness & Care in Mombasa, Kenya" (2010).  
*Independent Study Project (ISP) Collection*. Paper 820.  
[http://digitalcollections.sit.edu/isp\\_collection/820](http://digitalcollections.sit.edu/isp_collection/820)

This Unpublished Paper is brought to you for free and open access by the SIT Study Abroad at DigitalCollections@SIT. It has been accepted for inclusion in Independent Study Project (ISP) Collection by an authorized administrator of DigitalCollections@SIT. For more information, please contact [digitalcollections@sit.edu](mailto:digitalcollections@sit.edu).

**AN APPROACHING EPIDEMIC:  
EXPLORING DIABETES AWARENESS & CARE IN  
MOMBASA, KENYA**

BENJAMIN MARHAN DROPKIN  
HAMILTON COLLEGE

SCHOOL FOR INTERNATIONAL TRAINING  
KENYA: HEALTH AND COMMUNITY DEVELOPMENT  
SPRING 2010

ACADEMIC DIRECTORS: ODOCH PIDO AND JAMAL OMAR  
INDEPENDENT STUDY ADVISOR: MOHAMMED KARAMA

## **ACKNOWLEDGMENTS**

Many individuals helped make this independent study possible. I am deeply grateful to the twenty-five patients who graciously agreed to be interviewed and provided me with additional insight, contacts, and suggestions. These interviews would not have been possible without the help of Dr. Rishad, who allowed me to interview many of his patients and provided me with a number of additional medical contacts. I would like to thank Jamal Omar and Mohammed Karama for encouraging me to investigate diabetes in Kenya and for guiding me through the research process. I would like to thank my Kiswahili teachers Pete Obiero, Ann Tabu, and Geoffrey Maingi for teaching me to speak at a level sufficient to winning over patients with my often comical attempts at speaking the language. Finally, I would like to thank the students of SIT Kenya: Health and Community Development, Spring 2010 for helping to make this semester the most enlightening and enjoyable of my life.

## TABLE OF CONTENTS

ABSTRACT	3
INTRODUCTION	4
SETTING	10
METHODOLOGY	12
RESULTS & DISCUSSION	14
A. SURVEY RESULTS & ANALYSIS	14
B. BARRIERS TO IMPROVING DIABETES AWARENESS AT THE POPULATION LEVEL	20
C. SOLUTIONS TO IMPROVING DIABETES AWARENESS AT THE POPULATION LEVEL	24
D. BARRIERS TO IMPROVING DIABETES MANAGEMENT	30
E. SOLUTIONS TO IMPROVING DIABETES MANAGEMENT	34
CONCLUSION	37
RECOMMENDATIONS	39
APPENDICES	41
APPENDIX A: ORIGINAL PATIENT SURVEY	41
APPENDIX B: ADDITIONAL SURVEY QUESTIONS	42
APPENDIX C: GLOSSARY	42
BIBLIOGRAPHY	45

## **ABSTRACT**

The prevalence of Type II Diabetes Mellitus (T2DM) has become an increasing problem in virtually every city in the world due to lifestyle changes in recent decades that have greatly reduced healthy eating habits as well as exercise. Research has shown that T2DM is largely preventable through healthy lifestyle choices. It is crucial for every country's population to be aware of the risk factors and complications of diabetes so that they can take active steps to avoid the monumental physical and financial burden of the disease.

This study attempted to gauge diabetes awareness among patients living with diabetes and to explore barriers and solutions to both educating the public and improving diabetes management among current patients in Mombasa, Kenya. Interviews with 25 patients, 9 diabetes care providers, and a host of other professionals relating to diabetes were conducted over three weeks in April, 2010. Diabetes awareness among an educated patient sample was startlingly low. Barriers to public awareness and quality diabetes management of are numerous but feasible solutions appear to exist. To avoid the prevalence of diabetes reaching epidemic proportions, it is imperative that all citizens of Mombasa become aware of the serious health and financial risks posed by diabetes and make lifestyle changes accordingly.

## INTRODUCTION

Current estimates project the prevalence of diabetes mellitus (DM) world-wide to double from 175 million in 2000 to 353 million in 2030 (Yach, 2006:62). The majority of this increase is expected to occur within urban areas of developing countries like Kenya, where 305 million people are expected to have diabetes by 2030 (Yach, 2006:62). Ten years ago it was estimated that diabetes already accounted for 5.2% of deaths worldwide (Roglic, 2005:2130). DM needs to be addressed by all countries in order to control the enormous health and financial burdens that accompany the disease. There is no debate that the best way to achieve this goal is to avoid obesity through lifestyle choices. An important step in motivating people to make the necessary lifestyle changes is education about the risk factors and health risks of obesity and diabetes.

In recent years, most people in the developed world and an increasing number of those in developing countries have heard of diabetes. The number that has even a basic understanding of the disease, especially in developing countries, is radically lower. To fully understand the findings and conclusions of this study, it is important to have a basic understanding of diabetes. The same type of basic understanding, I would argue, that people everywhere would greatly benefit from having.

All forms of diabetes negatively affect the body's ability to maintain a relatively constant level of sugar in the blood. The most important hormone involved in this process is known as insulin, which is secreted by the pancreas and facilitates the uptake of sugar from the blood into the cells of the body (Mayo, 2010). Logically, if the pancreas is not able to produce insulin or if the insulin is not able to function properly then the body is not able to maintain a constant blood sugar level, which is harmful to the body. There are many types of diabetes but two types account for around 98% of all cases (WHO, 2009).

Type 1 Diabetes Mellitus (T1DM) is an autoimmune disease in which T lymphocytes (white blood cells) attack insulin-producing pancreatic beta cells and accounts for 5-10 percent of all diabetes cases (Herold, 2002:1692). T1DM used to be referred to as insulin-dependent diabetes mellitus (IDDM) or juvenile-onset diabetes because it tends to occur in children and adolescents and requires the use of synthetic insulin (NIH, 2007). The exact cause of T1DM has yet to be identified but risk factors may be genetic or environmental. No method of preventing T1DM has been identified (NIH, 2007). The prevalence of T1DM world-wide is increasing but not nearly at the rate that type II diabetes is (NIH, 2007).

Type 2 Diabetes Mellitus (T2DM) makes up 90% of all diabetes cases worldwide (WHO, 2009). T2DM refers to an impaired sensitivity to glucose in the cells of the body. At first, the pancreas increases insulin production to compensate for the decreased effectiveness but over time loses its ability to produce enough insulin (ADA, 2006:51). The result is consistently high blood sugar. T2DM results from an interaction between a genetic predisposition and behavioral and environmental risk factors, such as obesity and is associated with old age, physical inactivity, and race/ethnicity (Tuomilehto, 2001:1343). T2DM is usually diagnosed in overweight and obese adults, but diagnosis of adolescents is increasing around the world as well (NIH,2007).

The description of diabetes thus far might deter some people from unhealthy diets and encourage healthier lifestyles. Understanding the costs and complications of the disease, however, is a much stronger incentive to take steps to avoid the disease. Both T1DM and T2DM have been consistently shown to increase the risk of developing microvascular and macrovascular complications such as stroke, nephropathy (kidney failure), myocardial infarction (heart attack), retinopathy (loss of vision), and neuropathy (nerve damage which often leads to

numbness) (EAJM, 2006:397). The reason for these harmful and restrictive complications is that long-term hyperglycemia (high blood sugar) leads to damage of the vascular endothelium (a thin layer of cells that line blood vessels) as a result of oxidative stress (Brownless, 1995:223).

Encouragingly, all of these complications are reduced or avoided completely when patients are able to maintain a blood sugar level in the healthy range through a combination of diet, exercise, medications (often including insulin), and blood sugar management.

In a person without diabetes, the blood sugar level is normally regulated by the body to remain within a narrow range of 4.4 – 6.1 mmol/L (82 to 110 mg/dl) (ADA, 2006:51). For individuals with diabetes, all nine diabetes care providers I interviewed wanted their patients to maintain their blood sugar between 4 and 7 mmol/L. Three main monitoring techniques are used in diabetes management. In the first method, a patient places a drop of his or her blood onto a test strip which is coated with certain enzymes. The strip is then inserted into a glucometer, a hand held device which quickly delivers a measurement of the person's blood sugar at the time of testing. This test gives little indication of the quality of glycemic control over more than, at most, the last eight hours. The glucometer is an important tool for patients who correct their blood sugar with insulin injections. In a fasting blood sugar (FBS) test, a patient fasts for 8 hours before testing their blood sugar. This method gives a better picture of the patient's average blood sugar level. The best method for long term measurement is a laboratory test known as a Hemoglobin A1c (HbA1c). This method measures the average amount of glucose that has bound to the hemoglobin in the blood and provides a good measure of the patient's average blood sugar level over the previous three months (ADA, 2006:51).

In addition to the many harmful physical complications mentioned above, the financial burden associated with diabetes is alarmingly high. Diabetes costs in the U.S. were estimated to



be over US \$100 billion (KSH 75,000 billion) in 2000 (Zimet, 2000, 31). The average obese (BMI  $\geq$  30) American pays \$1,429 (in 2008 USD) more than the average American of normal weight (BMI < 25) (Flegal et. al, 2007:2028). These high costs are compounded in poorer countries like Kenya, where a greater lack of awareness often leads to later diagnoses where avoidable complications have already progressed (Davis, 1997:1435). In Mombasa, a diabetes consultation can cost up to 1500 ksh (Dr Rishad, 4/12/10). The vast majority of patients interviewed in this study reported the cost of medications as a major economic burden. The cost of blood sugar testing supplies inhibits most of the patients I interviewed from testing as often as they would like.

With an understanding of diabetes that includes these costs and complications individuals would be much more likely to make the difficult choices to eat healthier and get more exercise. Clearly the physical harm and financial burden that come with diabetes make the rapidly growing prevalence world-wide a major cause for concern. But what does this mean for Kenya? Practically every time I described my research project to anyone outside of the medical field I was met with the response “Diabetes in Kenya? Is that a problem there? I wouldn’t have expected that.” To which I now reply, “Remember asking me that in ten or fifteen years.”

In line with estimates of many African nations, the Kenyan ministry of health estimated the prevalence of Kenyans living with diabetes to be 1.2 million in 2007 and predicts the prevalence to rise to 1.5 million (4.5% of the population) by 2025 (Nation, 2007). The World Health Organization (WHO) estimated that in 2000 the prevalence of diabetes in Kenya ranked tenth of thirty four African countries, well behind countries like South Africa, Nigeria, and Algeria but ahead of countries including Uganda, Madagascar, and Mozambique. Also in 2000, the WHO rated Kenya’s healthcare system 140<sup>th</sup> out of 190 countries (WHO, 2000).

It seems fair to say that that Kenya is not unlike many other African countries in terms of diabetes and the care available. Cities all across Africa, and indeed the world, are facing many of the same problems as Mombasa. The barriers to improving awareness and care examined later in this paper should therefore be considered not as problems unique to Mombasa but as widespread problems that are plaguing or will plague many cities across the continent and world.

During my time in Mombasa I focused on two important questions: (1) ‘what do patients know about diabetes?’ and (2) ‘what can be done to improve public awareness and diabetes care?’ Given the time (three weeks) and resources available, I chose to interview 25 patients at four clinics, 9 diabetes care providers, 1 diabetes clinic director, 2 laboratory technicians, 3 pharmacists, 1 Chief (an appointed official who serves one of the 2500 ‘locations’ in Kenya) , 1 Senior Education Official, 1 teacher/administrator and 1 secondary school student. The range of professionals that I talked to is indicative of the many barriers to improvement that I found and explored. I originally chose to interview patients with diabetes instead of non-afflicted individuals in order to examine the factors associated with good glycemic control. When it became apparent that diabetes awareness among patients was so startlingly low, I did not see a need nor have the time to extend the interviews into the general public. It is obvious that the population as a whole needs to become educated about the risk factors and complications of diabetes and make lifestyle changes accordingly.

I chose to focus on diabetes in Mombasa because of my previous experiences related to diabetes in the past few years and because the problem has the potential to cripple the population and health care system if not addressed immediately. Indeed, in America we are already seeing a full third of the population classified as obese (WIN, 2010). Examining possible solutions intrigued me because of the enormous health-related, societal, and economic implications of the

challenge of diabetes. While communicable diseases like HIV have been the major challenge of the last thirty years, I believe non-communicable diseases like diabetes are going to be the major challenge of the next thirty. I chose to conduct this study in Mombasa primarily due to medical contacts made available to me by SIT Nairobi academic director, Jamal Omar.

Once it became clear that patient volume and reliable medical data were going to be much harder to come by, my goals for this project shifted from assessing the factors associated with good glycemic control to gauging diabetes awareness among patients and exploring the barriers and solutions to improving diabetes awareness and care in Mombasa.

## SETTING

All interviews and research presented in this study occurred in Mombasa, Kenya during April 2010. Mombasa is an island city of 730,000, predominantly Muslims and Christians, on Kenya's East coast surrounded by the Indian Ocean. Mombasa's economy is largely connected to the Kilindini Harbor, the only large harbor in Kenya. Tourism, oil production and cement production also contribute significantly to the economy. An important factor that probably contributes negatively to exercise rates is an average daily high temperature of 84 degrees Fahrenheit and high humidity (BBC, 2010). During my time in Mombasa it was the "rainy season," yet the daily high temperature hovered around 90 degrees Fahrenheit. According to a number of professional I talked to, significantly fewer patients come in when it is raining, so the daily rains may have decreased patient volume during my research period.

Each interview occurred in one of the following areas: Old Town, Kizingo, and Kibokoni. These three affluent districts contain numerous medical clinics and hospitals of varying quality, traditional/herbal healing options, and a very high number of chemists (pharmacies). The clinics where I conducted interviews charged consultation fees ranging from 250 ksh (3.33 USD) to 1500 ksh (20 USD) per visit. Almost all of the patients I surveyed reported the costs of their consultations and medications to be a significant financial burden. Nonetheless, the sample of patients I spoke with generally represents the middle-upper class of Mombasa. It is logical to hypothesize that this portion of the population knows more about diabetes than the lower classes because of better education and access to health care. My findings on diabetes awareness, therefore, are likely a best case scenario for the population as a whole at present.

Two recent societal changes are undoubtedly contributing to the increasing prevalence of diabetes and obesity in Mombasa. Like most cities in the world, the introduction of readily available, affordable, and unhealthy cooking materials and fast food options such as frying oil and chicken and chips has occurred within the last few decades. A fundamental shift in diet across Kenya and many other countries has resulted. Traditional dishes like mashed potatoes with beans and salt have been altered to include cream and fat or replaced altogether with dishes like fried chicken and chips (Dr. Gaman, 2/8/2010). Restaurants that serve cheap, unhealthy food are ubiquitous in Mombasa. One restaurant near Mombasa Hospital serves a very large plate of pilau (spiced, fried rice with chunks of meat) for just 50 ksh (.66 USD). Bottles of sugared soda are available on every street for 30 ksh (.40 USD). Ice cream and numerous fried foods such as samosas (fried meat pies) and mandazi (donuts) are available on practically every street in the city.

The second change, the introduction of tuk-tuks, seems to be less widespread but certainly not unique to Mombasa. Tuk-tuks are cheap, dangerous, and very maneuverable three-wheeled taxis that charge a fraction of the price of their four wheeled competition. Tuk-tuks only became widespread in Mombasa four years ago and have not yet become widespread in Nairobi (Interview #16, 4/21/10). Tuk-tuks have become a staple of transportation for Mombasa residents and have significantly reduced the need to walk. This is especially worrying in light of the fact that 'walking' was practically the only form of exercise reported during this study. Multiple health care providers mentioned tuk-tuks as a major obstacle to getting people to exercise more.

## METHODOLOGY

When I originally created the survey found in Appendix 1, my goal was to identify predicting factors associated with good diabetes management. From previous conversations with my advisors in Mombasa and Nairobi, I imagined that I would be able to survey a sample of around 100 patients with diabetes and conduct a brief statistical analysis on the results. When I arrived in Mombasa and went to the Aga Khan clinic for the first time, I quickly realized that this was not going to be feasible. In the United States, doctors specialize in endocrinology after medical school and can fill the majority of their schedules with cases related to diabetes. Doctors are incentivized to see a high patient volume in order to pay their bills and make a profit. Many end up seeing 25 patients with diabetes in a single day. All practices in the U.S. must keep accurate records, many of which are electronic, for insurance, liability, billing, and follow up care purposes.

I expected something similar in Mombasa but I found something very different. Many providers of diabetes care are general practitioners who run a diabetes clinic one morning or afternoon a week. Others, like Dr. Rishad, have specialized in another field (nephrology in his case) but see a handful of patients with diabetes each day. Medical records are rarely easily accessible and were never electronic in the settings where I conducted interviews. I quickly realized that collecting accurate diagnostic information such as blood pressure, weight at diagnosis, and HbA1c, was not going to be feasible given the time and resources allotted to me.

Despite this challenge, it became apparent on my first day that patients living with diabetes knew very little about the disease. I quickly modified the survey to contain questions about the cause of diabetes, source of patient knowledge, immediate reason for diagnosis, and care seeking behavior. These additional questions are listed under Appendix 2. Using this new

survey, I interviewed 10 patients at Dr. Rishad's Aga Khan Clinic, 4 patients at his Mombasa Hospital Clinic, 8 patients at Dr. Lema's Mombasa Hospital clinic, 2 patients at Dr. Bachani's Memon Clinic, and 1 patient at the SIT Mombasa office over the course of three weeks. In addition to the survey, I asked some of the patients about educating the population and ways to reduce the incidence of the disease.

In order to identify and address as many of the barriers to educating the population as possible, I began to interview a range of other people with jobs relating to diabetes. When one person would mention a seemingly legitimate barrier to educating the public, I sought out someone in that field to better understand the issue. For example, Dr. Lema mentioned that he believed education about diet and diabetes in schools was the best solution so I went and interviewed an official at the Ministry of Education and a teacher/administrator at a secondary school. Dr. Ashraf told me that many patients got frustrated with conventional medicine and transitioned to alternative medicine so I went to an herbal medicine clinic and interviewed an alternative medicine provider. In total I interviewed 6 medical doctors, 2 clinical officers, 1 herbal/alternative clinician, 2 laboratory technicians, 3 pharmacists, 1 diabetes clinic director, 1 chief, 1 senior education officer, 1 teacher/administrator, and 1 secondary school student. In each case, I prepared a list of specific questions and got permission from the individual to use the information they provided in my report.

The majority of patients I interviewed spoke English well enough that no translation help was needed. In the five cases where help was required, I was assisted by (3) Dr. Rishad's secretary, (1) a staff member at Dr. Lema's clinic, (1) another patient in the waiting room, and (1) a patient's daughter. I do not see the language barrier as a significant limitation to this study.

## RESULTS & DISCUSSION

### A. SURVEY RESULTS & ANALYSIS

#### I. DEMOGRAPHIC INFORMATION

	<b>Total</b>	<b>Men</b>	<b>Women</b>
<b>Sample Size</b>	25	12	13
<b>Average Age</b>	52	52	52
<b>BMI</b>	31.2	28.6	33.9
<b>Average Visits / Year</b>	17	10	22
<b>Average Years Since Diagnosis</b>	5.5	4.9	6.2

Table 1. Demographic Information of the Twenty-Six Patients Interviewed.

The sample tended to be well educated, living in urban areas within Mombasa, and practicing either Islam or Christianity. All but four of the respondents had completed at least secondary school. Economic status was not specially discussed but based on the consultation fees of the private practices where interviews were conducted it is safe to say that this sample was average to well above average financially. I expected that this portion of the population would be the most informed about diabetes based on access to health care, education level, and wealth. One patient even characterized type II diabetes as “a disease of the rich” (Interview #24, 4/28/10). We can therefore take the following results regarding awareness, testing habits, and care-seeking behavior as a best case scenario for Mombasa.

#### II. GENERAL DIABETES AWARENESS

The first step in convincing someone to make difficult lifestyle changes to avoid diabetes is convincing them that diabetes is avoidable. All people need to understand that type II diabetes is extremely preventable via lifestyle choices. People with normal BMIs (BMI<25) are much less



likely to develop type II diabetes than those who are overweight ( $BMI \geq 25$ ) or obese ( $BMI \geq 30$ ) (Knowler et. al, 2002:393). I asked 25 of the patients I interviewed what caused diabetes. Eleven said they had no idea whatsoever. Some interesting secondary comments came from this group. One patient said that “only God can know that” (Interview #6, 4/13/20). Two said that “[they] couldn’t know that because [they] were not doctors” (Interviews # 14 & #22). One patient told me that “like HIV, I believe the Europeans may have made diabetes so that they could sell their drugs” (Interview #23, 4/29/10).

Nine patients mentioned genetics or inheritance as the primary cause of diabetes. This issue is discussed in the next section. Two patients mentioned “stress” as the primary cause, one of which told me how this explanation accounted for the high rates of obesity in rural villages. He said that rural villagers have at least ten children, which is very stressful, and then naturally get diabetes. It is important to keep in mind that the people responding to this question have had diabetes for an average of five and half years and visit a doctor for care an average of seventeen times per year.

Only four out of the twenty five mentioned weight or lifestyle as a contributing factor to developing diabetes. This is probably the most concerning result of this study. Among a group of generally well educated, financially stable, people *who already have diabetes and see a provider* only 4 out of 25 mentioned lifestyle habits as a cause of diabetes. Two of these had attended nursing school. So among patients with no medical training, only 2 or 23 knew anything about the link between poor diet and exercise and developing diabetes.

### III. AWARENESS OF COMPLICATIONS

There is no question that people who spend more time and effort trying to maintain a constant and healthy blood sugar level if they knew of the terrible long term complications of diabetes: vision loss, kidney failure, heart problems, numbness of the hands and feet, delayed wound healing, and more (EAJM, 2006:397). When asked “what are some of the long term complications of diabetes?” most responses included one to three of these proven complications. Five patients said that they didn’t know of any complications of the disease. One patient said that “if you get diabetes, it means you already have TB and your joints are going to be destroyed” (Interview #16, 4/21/10). Educating patients about the many complications of diabetes is an important step in motivating them to control their blood sugars.

### IV. GLYCEMIC CONTROL AWARENESS

According the American Diabetes Association (ADA), the range of blood sugar that patients should aim for is between 4 and 7 mmol/L (ADA, 2006). Encouragingly, almost all of the patients who responded to this question gave ranges within 1 mmol/L of this range. Four said they had no idea and two gave very high target ranges: 6-10 mmol/L and 7-10 mmol.L, respectively. It is important that all patients know the range the ideal range to avoid short term symptoms and long term complications.

## V. BLOOD SUGAR MONITORING

<b>Location</b>	<b>Responses</b>	<b>Frequency of Testing</b>	<b>Responses</b>
Hospital/Clinic	5	During Visits	3
Home	13	At least Once per Day	7
Chemist	1	Weekly	4
Doesn't Check	1	Monthly	5

Table 2. Location and Frequency of Blood Sugar Testing in the Sample Group

Frequent blood sugar testing is a critical component of good diabetes management. Ideally, patients using insulin should check their blood sugar at least before each meal if not more frequently. Frequent testing can help patients avoid symptoms of both high and low blood sugar. If a patient is taking insulin, for example, a low blood sugar level would indicate to them that they should reduce the amount of insulin they are injecting. Not one patient described testing in this fashion. Most patients who test at home said they would test more frequently if the strips weren't so expensive. I visited three pharmacies and found that on average a package of 50 one-time-use strips cost 1,879 ksh. If patients were to test their sugar twice a day, this cost would become a monthly expense that few patients could afford. Many patients seem to check their blood sugar with a glucometer once a week or less. This is not a great way to monitor blood sugar because it only gives the patient a short term picture of their control. A number of patients said they only tested when they felt the symptoms of high blood sugar which may mean that they are often have blood sugar levels which are well above the desired range but not yet producing symptoms. One patient reported not being able to afford to test ever (Interview #17, 4/10).

## VI. DIET

The foods reported as being dietary staples did not vary significantly across the sample. The traditional Swahili diet includes rice, ugali, chipati, beans, fish, chicken, vegetables, fruits, etc. All of these answers were very common. Many patients said they had switched away from white flour products like white rice and white bread for the brown flour alternatives. Like any society, it is hard to imagine adults changing their diet after eating the same foods for their entire lives. The tendency to keep the same diet may be magnified in a place like Mombasa, where the standard diet is very narrow and few alternatives are both available and affordable. Patients need to be encouraged to reduce their portion sizes and use techniques to minimize the number of carbohydrates they are consuming. An easy technique mentioned by one patient is to let rice soak for an hour or so and remove the excess starch before cooking (Interview #10 4/10).

## VII. POST-DIAGNOSIS NUTRITIONAL COUNSELING

Almost every patient reported receiving advice about limiting portions, avoiding sugary and starchy foods, and opting for brown flour products over white flour products. Exceptions to this trend included several patients who said they had never been counseled on the topic. One 80 year-old gentleman, who was diagnosed in 1968, said that he had no idea which types of food to avoid or reduce (Interview #14, 4/10). Despite listing sugary and starchy foods, many patients were not able to mention many actual examples of these types of foods. Between nutritional counseling and posters or pamphlets illustrating actual examples of these types of foods patients could become better informed. One recently diagnosed patient said that he wanted nutritional counseling but had not received any from his doctor and did not know where to look (Interview #2, 4/12/20).

## VIII. EXERCISE

Fourteen patients described walking as their only form of exercise. Two added swimming. The remainder said that they didn't get any exercise anymore. A culture of physical activity needs to be promoted on the population level. This issue is addressed in Discussion Section C.

## IX. FAMILY HISTORY

It was very common for patients to have a family member afflicted with diabetes. Thirteen patients responded that they had a relative with diabetes while ten responded that they were the first in their family. There is no doubt that genetics plays a role in diabetes. The data from these responses alone, however, can tell us nothing concrete. It could be that patients with diabetes tend to come from families who can afford and choose to eat large portions of unhealthy foods and get very little exercise. Interestingly, three patients without a family history of diabetes reported that their spouse also had the disease. Nonetheless, the frequency of multiple members of the same family getting diabetes is contributing to the notion that diabetes caused by genetics. This is a very dangerous idea which is discussed in depth in the Discussion Section B.

## X. MEDICATIONS & AFFORDABILITY

Patients reported a wide range of medications including several types of insulin (lantas, humalog, etc.) and tablets (diamacrin, metformin, glucophage etc.). Only one patient reported an adverse effect of a medication. Medication availability is not an issue in Mombasa. Medication affordability, however, was described as a major economic burden by practically every patient. Three respondents said that their costs are not an issue at the moment because of employer

insurance policies but that they didn't know if they could afford the medications when they retired. Two patients said that they bought their medications from India at a much better price than is available in Mombasa.

## **B. BARRIERS TO IMPROVING DIABETES AWARENESS AT THE POPULATION LEVEL**

### **I. PRIMORDIAL EDUCATION**

The term 'primordial education' refers to teaching a subject starting on the first day a child attends school. When I first met with Dr. Lema, he told me that primordial education was the best chance we have at combating the current trends in obesity and diabetes (Dr. Lema, 4/14/10). While his conviction was strong, Dr. Lema didn't seem to know much about what, if anything was being taught to primary and secondary school students about diet, nutrition, and the complications of obesity. Looking for more information, Charles Nyaribo, a Senior Education Officer at the Ministry of Education reported that both private and public schools teach the same basic curriculum, which is designed by the Kenya Institute of Education (SIE) (Mr. Nyaribo, 4/22/10). He mentioned that students learned about health in a class called "home science" but referred me to a teacher at the Coast Girls Secondary School for more specific information.

Mrs. Thirika is the acting head of the Creative Arts department at Coast Girls. As for primary education, Mrs. Thirika informed me that "science and crafts used to be taught but has been removed from the curriculum," leaving no education related to diet or exercise at the primary school level. During the first day of Form I (9<sup>th</sup> grade), students choose one of four classes to take for the next four years: Home Science, Computer, Business, or Agriculture. After the completion of Form II, students have the option to transfer classes but *do not* have the option to transfer into Home Science. Currently, 40% of the Form I class at Coast Girls is enrolled in

Home Science (Mrs. Thirika, 4/23/10). Mrs. Thirika referred to the students who choose Home Science as “the lucky ones.”

Home Science teaches child care, home management, consumer education, clothing & textiles, health, and food and nutrition. Students in Forms I and II attend Home Science three times per week while students in Forms III and IV attend it four. Two lessons per year are taught on “Nutritional Deficiency Disorders,” during which the girls learn about hypertension, goiters, marasmus (a severe protein-energy malnutrition), and diabetes. Aside from occasional lectures on topics like HIV and TB, the information taught in Home Science is the only information about health or nutrition that students receive during their entire educations. I interviewed a secondary school student who had opted for Computer class in his secondary school and could not recall ever learning a thing about health or nutrition aside from a pamphlet distributed by a visiting head of school, which few students read (Mr. Khan, 4/22/10). Mrs. Thirika emphasized the importance of educating students about diabetes so that they would be able to go “go back and teach their relatives.” She also said that “the majority of students have a relative at home with diabetes” (Mrs. Thirika, 4/23/10).

Besides the obvious problem that the majority of students are not even enrolled in Home Science, many schools are cutting the course because of the expenses that come with it. Principals are arguing that the class is not affordable because of the high costs of supplies, which include cooking devices, utensils, materials, sewing machines, kitchens, fabric, etc. Students are already required to bring their own cooking supplies from home, but during examinations the schools must buy the supplies to standardize the exam (Mrs. Thirika, 4/23/10).

Outside of diet and health education, a Physical Education (P.E.) program is in place teaches students gymnastics, games, swimming, etc. Currently each class spends only forty minutes per week in P.E. (Mrs. Thirika, 4/23/10).

## II. AN “IT’S GENETIC” MENTALITY

Why asked “what do you think causes diabetes?” nine of the twenty-five patients responded “it’s genetic” or “it’s inherited.” It is well established that genetics can predispose certain people to obesity and diabetes (Weber, 2008:65). It is equally well established that lifestyle characteristics like diet and exercise also play a major role in determining who gets diabetes. Numerous “lifestyle intervention studies” have shown that diabetes can be prevented through changes in behaviors even among those containing strong genetic predispositions (Knowler, 2003:393).

An “it’s genetic” mentality is appealing to people because it removes any responsibility they might feel to eat well and exercise. One patient with a BMI of 53.8 (morbidly obese) told me that “they say it’s caused by lifestyle and too much weight, but I think it’s genetic” (Interview #13, 4/10). This patient’s sisters, brothers, and mother all had diabetes. Chances are that they were also all obese. Very few patients seem to realize that just because everyone in a family is obese and has diabetes, we do not have any more reason to believe that the disease is genetic than we do to believe that it is caused by obesity (Dr. Rishad 4/27/10). Unfortunately, choosing the “it’s genetic” mentality often leads patients to ignore advice about changing lifestyle habits and leads them down the painful road of diabetic complications.

It is also possible that the limited diabetes awareness of providers is contributing to the “it’s genetic” mentality. According to Dr. Okanga, a former branch chairman of the Kenya



Diabetes Association, most doctors only learn about diabetes for a semester or two in medical school and then pick up bits of information during their internships. He thinks it is likely that many doctors providing diabetes care are unsure about the etiology of the disease themselves and may be giving patients the easily accepted answer that the disease is genetic (Dr. Okanga, 4/28/10).

### III. GOVERNMENT PRIORITIES

The rates of non-communicable diseases like diabetes are increasing all over the world. Developing countries like Kenya are facing a “double burden” of both non-communicable and communicable diseases that they must deal with (Boutayeb, 2001:191). Right now the Kenyan government is more concerned with combating HIV, malaria and tuberculosis than with diabetes because the former are thought of as the main medical problems in the country. Bakari Mohammed, Chief of Old Town, told me that health-related talks do occur at the bimonthly Chief’s Barazas but that they are almost always related to TB, malaria, and clean drinking water. He was very vague on the frequency of talks related to nutrition or diabetes, but did say that they do exist “because diabetes is a very big problem for [their] people” (B. Mohammed, 4/23/10). It’s very hard to say that the money spent on HIV, malaria, and TB education would be better spent on diabetes. It is fair to say that health care costs over the few decades could be radically reduced if more money was spent on nutrition education now.

### IV. GOVERNMENT FUNCTIONING

Like all sectors of the government, the health sector is prone to abuse and misuse of resources. Several people I interviewed expressed their pessimism about the government being

able to achieve anything to benefit the health of the population. One woman who spends most of her time helping sick people in her community told me that the government corruption she's witnessed makes her "wish she was never born" (N. Mohammed, 4/20/10). Dr. Okanga told me about a serious mismanagement of the funds given to the Kenyan Diabetes Association by the International Diabetes Foundation (IDF) in 2008 (Dr. Okanga, 4/28/10). He described the corruption as crippling to the progress the KDA was making with their education outreach programs.

#### V. FUNDING FOR PUBLIC EDUCATION

Like any country, but particularly developing ones, funding for anything is hard to come by. Often times when a health-related issue can lead to profits for a corporation, that corporation will invest in educating the population on the topic through things like pamphlets and TV commercials. While drug companies do have an incentive to educate patients about drugs that can help to control diabetes there is no incentive for those companies to encourage lifestyle changes that could help people avoid diabetes in the first place (Dr. Ashraf, 4/16/10). Organizations seeking private funding must compete with the same prevalent and highly publicized diseases which are absorbing much of the government funding.

### **C. SOLUTIONS TO IMPROVING DIABETES AWARENESS AT THE POPULATION LEVEL**

#### I. ORGANIZED OUTREACH PROGRAMS

Diabetes Kenya (formerly the Kenyan Diabetes Association) and the World Diabetes Federation are playing important roles in educating the population and identifying individuals with diabetes. Diabetes Kenya's strategy involves establishing diabetes clinics in locations all

over Kenya including places like Lamu, Melindi, and Kilifi. Diabetes Educators at these clinics train others in the surrounding areas to go back to their communities and conduct talks about diabetes at mosques, churches, and schools and look for people experiencing the symptoms of diabetes. Dr. Okanga said that the system was working very well before IDF funds were poorly managed in 2008. After a rough patch with limited funding, the project now seems to be improving the situation once again (Dr. Okanga, 4/28/10). This type of widespread outreach has the potential to prevent people from getting diabetes and to help those with diabetes get care as soon as possible.

The World Diabetes Foundation and the Diabetes Information and Management centre conducted the Diabetes Education Program (WDF04-085) from 2005-2009 hoping to “improve the understanding of diabetes among health care staff and create public awareness on prevention, control and risk factors” (WDF, 2009). The program created 45 main clinics at district hospitals, 109 mini-clinics at dispensaries and educated 682 doctors, 2,091 nurses, 813 paramedics, 380 dieticians, and 4,023 lay educators. The program also screened 118,767 people across the country and found that 12,627 (over 10%) had diabetes, 1,834 of which were unaware they had diabetes. One-day educational meetings were held within religious and corporate settings, educating a total of 661,638 people.

The WDF is now involved in two additional projects launched in 2008 and 2009 targeting diabetic foot care (WDF07-302) and Diabetes Care in the slums of Nairobi (WDF08-399), respectively (WDF, 2009). The results of the initial program show what an important and effective role an international organization like the WDF can play in a developing country like Kenya. As encouraging as these results are, it must also be kept in mind that Kenya is a country

of 39 million people (CIA, 2010). Educating 660 thousand people, therefore, is a small but important first step towards the goal of near complete population awareness.

## II. PRIMORDIAL EDUCATION

Graduating every student from secondary school with a solid understanding of basic nutrition, diet, exercise, the risks of obesity and the signs and symptoms of diabetes is a realistic and important goal. Requiring students to take and adequately funding the Home Science class would be one way to reach this goal. Because increasing government funding is such a difficult task, other solutions need to be explored. School-wide lectures regarding communicable disease like HIV are already commonplace. Mrs. Thirika saw no reason why diabetes-related lectures of the same kind could not be utilized. She also thought that diabetes education could easily be added to current topics addressed by each school's Guidance and Counseling Committee. In addition to the long term benefits of primordial education, students are able to pass on the things they learn to their families and friends who may have diabetes or be in a high-risk group.

The longer a person with diabetes remains undiagnosed the worse their prognosis is becomes. Knowing the symptoms of diabetes is important for making diagnoses as early as possible. Towards the middle of my research period I began asking patients about how they were diagnosed. One woman told me that she had been urinating frequently and was very thirsty so she went to get checked for diabetes. When I asked her how she had known that those were signs of diabetes she told me that she had learned about diabetes in secondary 25 years ago. She is now 38 years old. Knowing the signs of diabetes probably sped her recovery and significantly increased her life expectancy.

In 2009, The Ministry of Health in the United Arab Emirates (UAE) has launched a three year campaign to educate students, parents, and teachers about diabetes by holding frequent workshops held at primary and secondary schools in UAE (DN, 2010). Obviously more funding for this type of endeavor is available in the UAE than in Kenya at the moment. Nonetheless, this campaign could serve as an excellent model for Kenya.

## II. DIABETES CAMPS

“Diabetes Camps” are days where free head-to-toe checkups are provided at certain hospitals in Mombasa. Alice Halwenge, the director of the Mombasa Hospital Diabetes Clinic, said that the hospital runs two camps per year and that hundreds of people come to each. At these camps, patients have their blood sugar checked and receive free counseling on diet and exercise from nurses. Chief Mohammed also mentioned the camps as one of the major ways that people in his district become educated about diabetes (B. Mohammed, 4/23/10). Increasing the frequency of diabetes camps is a desirable goal that is contingent on funding available to the participating hospitals

## III. CHIEF’S BARAZAS

Chief’s barazas were mentioned in three interviews as a way to educate the public so I went to talk with a Chief in Old Town. A baraza is a large gathering where a district’s population meets at a local school or mosque (B. Mohammed, 4/23/10). Most districts conduct around twenty-four barazas each year. These gatherings often include health-related lectures but it seems that currently the majority are about TB, malaria, and clean drinking water. Adding lectures about nutrition and diabetes to this slate seems like a relatively cheap and feasible way to educate

the population. Mr. Mohammed supported the idea in light of the increasing burden of diabetes in his community. Finding an engaging speaker to lecture on these topics is a potential problem (B. Mohammed, 4/23/10).

#### IV. WAITING ROOM EDUCATION

Most of the medical clinics I visited in Mombasa functioned on a first come, first serve basis. This often meant many patients waiting idly over an hour to see a doctor. This would be a great time to educate patients about nutrition and diabetes. I was advised not to visit Coast General Hospital because of protocol issues, but several patients told me about diabetes educators who lecture in the waiting areas there. One Clinic I went to is planning to install a television in the waiting room that will show educational videos on repeat (Dr. Bachani, 4/21/10). Keeping in mind the high start up cost of buying a television and video player, this method seems like a great way to get information to patients while they are not busy.

#### V. EDUCATING ALL PATIENTS

Physicians in all specialties should be comfortable and knowledgeable enough to talk to any of their patients about basic nutrition, the risks of obesity, and signs and symptoms of diabetes. Whether seeing a patient for hypertension or a knee replacement, if a doctor encounters an obese patient they should at least encourage that patient to lose weight and investigate the signs of diabetes.

## VI. WORLD DIABETES DAY

World Diabetes Day occurs two weeks after Diabetes Awareness Month (November) concludes, on December 14<sup>th</sup>. Since 2004, the World Diabetes Foundation (WDF) has organized the Global Diabetes Walk. Last year's walk included over 305,000 participants in 343 registered walks in 52 countries including Kenya. Mombasa was a host city for one of these organized walks. This is a great event because it is highly publicized and brings together the diabetes community in a city for a day where information and knowledge about management and prevention can be shared (WDF, 2009).

## VII. PROMOTING PHYSICAL ACTIVITY

A landmark study by Knowler et al. randomly assigned 3,234 individuals at high risk for developing diabetes to one of three interventions: lifestyle changes, the drug metformin, and information about diet and exercise alone (Knowler et. al, 2002:393). "Lifestyle changes" usually meant getting 30 minute of exercise (usually walking) 5 days a week and decreasing calorie consumption. The study found that those in the lifestyle intervention group reduced their risk of getting type II diabetes by 58 percent, while those in the metofromin group reduced their chances by 31 percent (Knowler et al, 2002:393). Thirty minutes of walking per day may have once been the norm in Mombasa, but since the introduction of cheap and available transportation including matatus and tuk-tuks, it certainly not any more.

The best way to increase the physical activity of a population is to develop healthy exercise habits among children that can be sustained throughout life. Increasing the amount of P.E. in schools is a critical first step to achieving this goal. The standard forty minutes of PE per school week is insufficient by any measure (Mrs. Thirika, 4/23/10). Enjoyment of physical

activity is becoming increasingly important as less and less exercise is required on a daily basis. Additional efforts need to be made to encourage adults to get exercise each day. The most realistic forms of exercise in Mombasa are walking and swimming. Although the traditional Muslim dress could be a problem for some women, I saw many Muslim women walking and running in burkas on my own morning jogs. In the U.S. the National Football League (NFL) has launched a widespread campaign called Pay 60: the NFL movement for an active generation. This initiative encourages communities to facilitate children playing actively for at least an hour a day (NFL, 2010). Again, funding would be much harder to garner in Kenya, but the method of prevention is a good one.

#### **D. BARRIERS TO IMPROVING DIABETES MANAGEMENT**

There is no question that prevention through lifestyle choices is the best way to combat the burgeoning problem of diabetes in Mombasa. For those who have or will soon have diabetes, it is very important that quality medical care be available to help patients manage their diabetes in order to avoid symptoms and complications of the disease. There are a surprising number of barriers to improving diabetes management in Mombasa, many of which I have not seen during my experience in the United States.

##### **I. PATIENT COMMITMENT**

Glycemic control and the resulting symptoms and complications are largely products of an individual's effort to manage their diabetes. Patient commitment to the management of diabetes appears to be a very serious problem in Mombasa. Practically every doctor I interviewed commented on this problem. Dr. Ashraf said that one of his biggest challenges



within diabetes care is convincing patients to follow up (Dr. Ashraf, 4/16/10). Dr. Lema told me that “patients with decent or good control don’t come in for appointments” (Dr. Lema, 4/14/10). Mrs. Halwenge told me that “many patients will not come in unless they’ve had recent symptoms” (Mrs. Halwenge, 4/14/10).

In response to these comments, I began asking the patients I interviewed why they came in for diabetes care. It was not uncommon for a patient to respond that they came only when they experienced symptoms. While the average number of visits per year displayed in table 1 is not bad, this average is somewhat deceiving because a handful of very committed patients increased the overall average significantly. Alarming, it seems that the weather can be a major factor in patients’ decisions to visit the clinic. Many diabetes clinics are only open one day per week. If it happens to be raining that day, many patients will choose not to visit the clinic that week (Mrs. Halwenge, 4/14/10).

It would make sense that commitment to diabetes management would be strongly linked to an understanding of the benefits of good control and of the possible long term complications of the disease. This is certainly an area for future study which could be important in garnering outside funding for diabetes education at the population level.

## II. QUALITY OF CARE PROVIDED

There is a great deal of room for improvement in the diabetes care provided in Mombasa. Diabetes education for doctors in Kenya is often limited to a semester or two in Medical School without much focus on the etiology of the disease (Dr. Okanga, 4/28/10). Clinical Officers receive limited training about diabetes that is largely limited to memorizing the symptoms and treatments of the disease (Dr. Collins, 4/16/10). Alternative medicine providers are usually

trained for one year by their employer and learn little more than the company's supplement and dosing recommendations for diabetes (M. Njoki, 4/19/10).

The product of this limited training is that many providers are not in a position to appropriately counsel patients about diabetes management. As mentioned earlier, the HbA1C is the best method to monitor a patient's average blood sugar control. When I interviewed Dr. Ali, a Clinical Officer, he was not familiar with the HbA1c or any other method of blood sugar monitoring besides the glucometer. This is a major cause for concern. If providers are not familiar with these basic testing methods then their patients have little chance of becoming educated. Providers with very limited training may be contributing to the perception that diabetes is solely a genetic disorder because of their lack of understanding (Dr. Okanaga, 4/28/10). This lack of understanding may also be leading some providers to use prescription medications as their only form of care because they are not in a position to educate.

Another major problem among providers is their unwillingness to encourage lifestyle changes (Dr. Ashraf, 4/16/10). They often feel that patients will not be able to make lifestyle changes, rendering their efforts useless. Financially speaking, doctors profit from patients returning for care often. It is possible that some doctors recognize this incentive and would rather see a patient monthly to renew their prescriptions than to help that patient avoid diabetes in the first place through lifestyle changes (Dr. Ashraf, 4/16/10).

### III. IDEAS ABOUT LOSING WEIGHT

Historically, a high weight has been considered a sign of affluence and success in much of Kenya (Kerns, 2006). While this trend may be slowly changing, partially due to western influence, it remains a serious problem for diabetes management. When told to lose weight,

many people are concerned about what their friends will think of them (Dr. Lema, 4/14/10).

From informal discussions I've had it also seems that very few Kenyans understand the association between BMI and life expectancy. A 2009 study found that BMI was a strong predictor of overall mortality both above and below the optimum range of 22.5–25. They found that at 30–35, median survival is reduced by 2–4 years and at 40–45 it is reduced by 8–10 years (Lancet, 2009:1083). One man who told me about his struggles to *gain* weight was shocked when I told him that thinner individuals tend to live longer than overweight and obese people.

#### IV. MANAGEMENT COSTS

Few people without diabetes understand how expensive good diabetes management is. At the locations where I conducted interviews, consultation fees ranged from 250 ksh to 1500 ksh per visit. The cost of medications is a major burden for virtually all patients paying out of pocket. One woman told me that she spends 6000 ksh on diabetes medications each month (Interview #8, 4/13/10). The monthly fee for the herbal supplements given to patients with diabetes at the Kamirithu Herbal Clinic is 3500 ksh (M. Njoki, 4/19/10). At the three pharmacies I visited the average price for a hand held glucometer was 3,608 ksh and the average cost of a package of 50 testing strips was 1,879 ksh. It seems that the majority of patients cannot afford to have a glucometer at home and therefore only check their blood sugar at their clinic or at a chemist. At the laboratory I visited at the Memon clinic, which had the least expensive consultation fee of the clinics I visited, it cost 150 ksh to test using a hand held glucometer and 700 ksh for an HbA1c. These costs are a major barrier to patients controlling their blood sugar. One patient at the Memon clinic told me that she stopped taking medications and never tests her blood sugar because she cannot afford to do so (Interview #12, 4/10).

## V. ALTERNATIVE MEDICINE

It is very hard to make general comments about the role of alternative medicine in diabetes care. There are a number of herbal and alternative medicine options for diabetes care in Mombasa. It is difficult to comment on the quality of care provided because studies measuring the efficacy of these treatments, which can differ greatly between providers, are extremely limited. According to Dr. Ashraf, many patients get frustrated with the slow progress they are making with conventional care and look for a quick fix through alternative medicine. Misinformation given to patients by families and friends about the effectiveness of these treatments also seems to be a contributing factor (Dr. Ashraf, 4/16/10).

Mercy Njoki, an alternative medicine provider at the Kamirithu Clinic, said that patients often come for supplements after they don't get well via conventional care. The providers working at the many clinics owned by Kamirithu are trained for one year at the company headquarters in Nairobi. All patients with diabetes who come to Karmirithu are given a supplement called "Kisukari." Ms. Njoki knew nothing about Kisukari besides that it was the supplement given for diabetes. According to Ms. Njoki, the supplements are used until the patient's blood sugar is back into a range of 3.5-6mmol/L, which usually takes between 6 and 12 months. Interestingly, if a patient can't get back into the normal range, they are advised to go to a conventional office and get insulin (Ms. Njoki, 4/19/10)

## E. SOLUTIONS TO IMPROVING DIABETES MANAGEMENT

### I. IMPROVING DIABETES AWARENESS

Literally every professional I interviewed felt that education was the best way to curb the trend in diabetes. For individuals already afflicted with the disease, education is still vitally

important both to help with food choices and to motivate patients to manage their diabetes as well as they can. As described at length in Discussion Sections B and C above, the barriers and feasible solutions to educating the public are numerous and complex. At the patient level, it is imperative that providers be able to raise the awareness of their patients, who will in turn be able to spread the word to others in their communities.

## II. MEDICAL EDUCATION

Medical education of all types of providers must include a thorough understanding of the etiology of diabetes and the current evidence that illustrates the importance of lifestyle choices in preventing the disease. Memorizing the symptoms and treatments for diabetes is insufficient knowledge if empowering patients is the goal. All providers must be in a position to educate their patients and answer their questions about diabetes managing. I would predict that as the rates of diabetes continue their upward climb, medical schools will increase their focus on diabetes out of necessity.

## III. EDUCATING CURRENT PROVIDERS

While education at the medical school level is very important for long term care, educating current providers is very important for care in the near future. The Kenya Medical Association (KMA) has introduced a Continuing Professional Development (CPD) program that is analogous to America's Continuing Medical Education (CME) system. Doctors are required to earn a certain number of points by doing things like attending conferences abroad or weekly lectures in Kenya in order to renew their licenses (Dr. Rishad, 4/26/10). Currently there are very few diabetes-specific lectures partially due to the government's primary medical concerns:

malaria and tuberculosis. Increasing the attention to diabetes could be an important way to increase knowledge among providers in the coming years.

## CONCLUSION

The prevalence of type II diabetes has been rapidly increasing across the globe in recent years. Lifestyle changes over the last decade have led to a diet of decreasing nutritional value and greatly reduced amounts of exercise. As a result, diabetes has transitioned from being a disease of the rich to a disease of the masses. Every person interviewed in this study reported that diabetes is a major problem for the population of Mombasa.

Preventative measures are the only way to keep the prevalence of diabetes in Mombasa from reaching epidemic proportions. The first step in prevention is educating people about the lifestyle choices that can prevent obesity, diabetes, and the harmful and financially draining complications associated with each. Alarmingly, only four of the twenty five patients interviewed in this study knew of the relationship between obesity and diabetes. Eleven had no idea what caused diabetes and nine thought diabetes was a strictly genetic disorder. This result is very bad sign for the near future of diabetes in Mombasa. If people don't understand that they can avoid the monumental physical and financial costs of diabetes through diet and exercise they are very unlikely to make difficult but important lifestyle changes.

Many barriers to raising diabetes awareness at the population exist. A severe lack of primary and secondary education, a prevalent "it's genetic" mentality, other important health burdens receiving the majority of government funding, and the government's inability to use funding efficiently and effectively are all major challenges. National and international organizations like Diabetes Kenya and the World Diabetes Foundation have begun efforts to educate the population within the last five years. An increased and sustained effort is undoubtedly required but may prove difficult to fund. Other ways to significantly increase public

awareness include primordial education, diabetes camps, chief's barazas, waiting room education, and the promotion of physical activity.

For those already living with diabetes, as well as the many who will become afflicted in the near future, several important barriers to maximizing diabetes management exist. These barriers include a lack of patient commitment (largely a product of not understanding the possible complications of poor management), the quality of diabetes care provided, negative ideas about losing weight among patients, and high the high costs of care and blood sugar testing supplies. To manage diabetes effectively, patients must be empowered through education about the multi faceted approach to managing the disease. For this to happen, it is imperative that a thorough understanding of diabetes is fostered among all current and future providers of medical care in Kenya.

Efforts at raising diabetes awareness now have the potential to save Kenya an immense amount of suffering and health-related costs over the coming decades. Every country should fear looking back in 2020 or 2030, with epidemic proportions of obesity and diabetes on their hands, and wishing they had invested more in prevention strategies. Based on the current status of diabetes awareness and care in Mombasa, it is hard to believe that this will not be the case.



## RECOMMENDATIONS

### A. EXPANSION OF THIS STUDY

With a longer research period I would have expanded my research in three ways. First, I would have liked to interview more patients, perhaps one hundred, across a more diverse socio-economic range than my sample provided in this study. To do this, I would have attempted to gain permission to conduct interviews at Coast General Hospital, the largest government-run hospital in Mombasa. Coast General has a large diabetes clinic where it seems that the patient volume is much higher and the financial stability of the patients is much lower than in the areas where I conducted interviews.

Next, I would have tried to further explore the role of the large organizations like the WDF and the challenges they face in a developing country like Kenya. Many issues are currently preventing the Kenyan government from functioning efficiently and effectively. International organizations may prove more effective at than government organizations at raising diabetes awareness in countries facing these types of challenges.

As I've stressed in this report, my sample group was a best case scenario of diabetes awareness and still knew extremely little about the etiology or management of the disease. Therefore my final expansion, given several months to work on the project, would have been to extend interviews into the general public to get a sense of diabetes awareness among those without the disease.

### B. FUTURE STUDY

Future studies related to diabetes in Mombasa should begin by trying to gain permission to conduct interviews at Coast General in order to maximize access to patients with diabetes. It

would be very interesting to reexamine patients' understanding of the etiology of diabetes and methods of prevention in the same locations as this study in five or ten years. In the near future, it would be very interesting to examine Kenyans' opinions about the relationships between BMI, health, and life expectancy. An understanding of these relationships is critical for people to make tough lifestyle decisions like eating healthier and exercising more.

Finally, it would be interesting to look into the pharmaceutical market in Mombasa. It appears that with a low barrier to entry, high profit margins, and ever increasing demand, more and more retail and wholesale pharmacies are opening in Mombasa. With seemingly little regulation or accountability, pharmacies often offer prescription drugs without prescriptions and may be selling counterfeit products. Based on the large price discrepancies I observed and the lack of any legitimate price-comparison method it seems that patients are paying radically different prices for the same medications.

## APPENDICES

### APPENDIX A: ORIGINAL PATIENT SURVEY

- I. Diagnostic Info:
  1. Age: \_\_\_\_\_
  2. Height: \_\_\_\_\_
  3. Weight: \_\_\_\_\_
  4. Weight when Diagnosed: \_\_\_\_\_
  5. B.P. \_\_\_\_\_
  6. Diabetes: \_\_\_\_\_
  7. A1C: \_\_\_\_\_
  8. Provider Rating: Terrible – Poor – Acceptable – Good - Excellent
  
- II. Lifestyle
  1. Education: primary \_\_\_\_\_ secondary \_\_\_\_\_ university \_\_\_\_\_
  2. Modes of Transportation: \_\_\_\_\_
  3. Exercise: \_\_\_\_\_
  4. Idea body type: diagram
  5. Which body type will live the longest?
  6. Where do you live? Rural – urban – slums
  7. Religious Orientation? \_\_\_\_\_
  8. Are you comfortable receiving injections?
  
- III. Diet
  1. Staple Foods: \_\_\_\_\_
  2. Importance of Price in food choices: \_\_\_\_\_
  3. Are there any foods that you were advised to stop eating or reduce quantities of after you were diagnosed with diabetes? \_\_\_\_\_
  4. Are there any foods that you have avoided since being diagnosed with diabetes?  
\_\_\_\_\_
  
- IV. Awareness
  1. When were you diagnosed / how long have you had diabetes? \_\_\_\_\_
  2. Has anyone else in your family had diabetes? \_\_\_\_\_
  3. What are some common complications of diabetes? \_\_\_\_\_
  4. What is good glycemetic control?
  5. Which foods increase your blood sugar the most? \_\_\_\_\_
  6. What are some causes of poor glycemetic control? \_\_\_\_\_
  7. How happy are you with your blood glucose levels on a scale of 1 (very unhappy) to 5 (very happy)? \_\_\_\_\_

## V. Healthcare

1. Are the drugs your doctor prescribes: available \_\_\_\_\_ affordable \_\_\_\_\_
2. Which type of medications do you take? \_\_\_\_\_
3. Do you think these drugs cause you any problems?
4. Have you ever failed to take your drugs?
  - i. If so, why?
5. How many times per year do you see your diabetes care provider? \_\_\_\_\_
6. Type of provider seen: consultant – GP – counselor
7. How do you check your blood sugar? \_\_\_\_\_
8. How often do you check your blood sugar? \_\_\_\_\_
9. Who do you go to when you get sick? \_\_\_\_\_
10. Do you seek healthcare anywhere besides the medical office?  
\_\_\_\_\_
11. Do you use any traditional treatments in addition to the drugs your doctor prescribes and which? \_\_\_\_\_
12. What are some of the symptoms you have experienced?  
\_\_\_\_\_

**APPENDIX B: ADDITIONAL SURVEY QUESTIONS**

1. What causes diabetes?
2. Where have you learned about diabetes?
3. How were you originally diagnosed with diabetes?
4. How could the population become more educated about diabetes?

**APPENDIX C: GLOSSARY**

**Autoimmune-** of, relating to, or caused by antibodies or T cells that attack molecules, cells, or tissues of the organism producing them.

**BG-** blood glucose- the amount of glucose (sugar) present in the blood of a human or animal. Normally, in mammals the body maintains the blood glucose level at a reference range between about 3.6 and 5.8 mM (mmol/L). It is tightly regulated as a part of metabolic homeostasis.

**BMI-** body mass index- a measure of body fat that is the ratio of the weight of the body in kilograms to the square of its height in meters <a *body mass index* in adults of 25 to 29.9 is considered an indication of overweight, and 30 or more an indication of obesity.

**Etiology-** the study of causation or origination.

**Glucose-** a sugar that can be linked to form carbohydrates and that serves as a primary source of energy.

**Glycemic Control-** a medical term referring to the typical levels of blood sugar (glucose) in a person with diabetes mellitus. Much evidence suggests that many of the long-term complications of diabetes, especially the microvascular complications, result from many years of hyperglycemia (elevated levels of glucose in the blood).

**HbA1C-** a stable glycoprotein formed when glucose binds to hemoglobin A in the blood ; *also* : a test that measures the level of hemoglobin A1c in the blood as a means of determining the average blood sugar concentrations for the preceding two to three months.

**Hyperglycemia-** an excess of sugar in the blood.

**Hypertension-** abnormally high arterial blood pressure that is usually indicated by an adult systolic blood pressure of 140 mm Hg or greater or a diastolic blood pressure of 90 mm Hg or greater.

**Macrovascular-** Pertaining to the macrovasculature, the portion of the vasculature of the body comprising the larger vessels, those with an internal diameter of more than 100 microns.

**Marasmus-** a form of severe protein-energy malnutrition characterized by energy deficiency.

**Microvascular-** of, relating to, or constituting the part of the circulatory system made up of minute vessels (as venules or capillaries) that average less than 0.3 millimeters in diameter.

**Nephropathy-** an abnormal state of the kidney; *especially* : one associated with or secondary to some other pathological process.

**Neuropathy-** an abnormal and usually degenerative state of the nervous system or nerves.

**Oxidative Stress-** physiological stress on the body that is caused by the cumulative damage done by free radicals inadequately neutralized by antioxidants and that is held to be associated with aging.

**Prevalence-** the percentage of a population that is affected with a particular disease at a given time.

**Retinopathy-** any of various non-inflammatory disorders of the retina including some that cause blindness.

**T lymphocytes-** a group of white blood cells that play a central role in cell-mediated immunity.

**Vascular Endothelium-** the endothelium is the thin layer of cells that line the interior surface of blood vessels, forming an interface between circulating blood in the lumen and the rest of the vessel wall. Vascular refers to a relation to blood vessels.

## BIBLIOGRAPHY

- A. Boutayeb. "The double burden of communicable and non-communicable diseases in developing countries." *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100 (2006):191-199.
- Alice Halwenge Interview, Director of the Mombasa Hospital Diabetes Clinic, April 14<sup>th</sup>, 2010.
- American Diabetes Association. "Standards of Medical Care-Table 6 and Table 7, Correlation between A1C level and Mean Plasma Glucose Levels on Multiple Testing over 2-3 months." *Diabetes Care* 29 (Jan, 2006): Supplement 1 Pages 51-580.
- Anonymous Patient Surveys #1-25, April 12<sup>th</sup>-28<sup>th</sup>, 2010. Mombasa, Kenya.
- Bakari Mohammed Interview, April 23<sup>rd</sup>, 2010. Chief of Old Town.
- British Broadcasting Company. "Average Conditions in Mombasa, Kenya." May 4<sup>th</sup>, 2010. <[http://www.bbc.co.uk/weather/world/city\\_guides/results.shtml?tt=TT000290](http://www.bbc.co.uk/weather/world/city_guides/results.shtml?tt=TT000290)>
- Brownless, M. "Advanced protein glycosylation in diabetes and aging." *Annw. Rev. Med.* 46 (1995): 223-234.
- Central Intelligence Agency. "World Fact book: Kenya." May 4<sup>th</sup>, 2010 <<https://www.cia.gov/library/publications/the-world-factbook/geos/ke.html>>
- Charles Nyaribo Interview, April 22<sup>nd</sup>, 2010. Ministry of Education.
- Davis, M. et al. "Effect of age of diagnosis on diabetic tissue damage during the first six years of NIDDM." *Diabetes Care* 20 (1997): 1435-1441.
- Diabetes News. "UAE launches diabetes awareness campaign in schools." April 14<sup>th</sup>, 2010. <<http://blog.taragana.com/health/diabetes/2010/04/14/uae-launches-diabetes-awareness-campaign-in-schools-1816/>>
- Dr. Ashraf Interview, April 16<sup>th</sup>, 2010. Ashraf Clinic at Posta.
- Dr. Bachani Interview, Memon Clinic, April 21<sup>st</sup>, 2010.
- Dr. Gaman Interview, February 8<sup>th</sup>, 2010. Nairobi, Kenya.
- Dr. Lema Interview, April 14<sup>th</sup>, 2010. Mombasa Hospital
- Dr. Okanga, April 28<sup>th</sup>, 2010. Aga Khan Doctor's Plaza.

Dr. Rishad Interview, April 26<sup>th</sup>, 2010. Mombasa Hospital.

Dr. Rishad Interview, April 27, 2010. Mombasa Hospital.

Flegal KM, Graubard BI, Williamson DF, et al. "Cause-Specific Excess Deaths Associated With Underweight, Overweight, and Obesity." *Journal of the American Medical Association*. 298 (2007): 2028–2037.

Herold, K. "Anti-CD3 Monoclonal Antibody in New-Onset Type 1 Diabetes Mellitus." *The New England Journal of Medicine* 346 (May, 2002): 1692-1698.

Kerns, C. "Smaller World, Smaller Bodies: The Shrinking Nature of the Globalization Beauty Ideal." SIT Fall 2006.

Knowler et. al. "Reduction in the Incidence of Type II diabetes with Lifestyle Intervention or Metformin: Diabetes Prevention Program Research Group." *New England Journal of Medicine* 346 (February, 2002: 393-403.

Mayo Clinic Staff. "Diabetes Treatment: using insulin to manage your blood sugar." <http://www.mayoclinic.com/health/diabetes-treatment/DA00010> Visited: March 24, 2010.

MedlinePlus Medical Dictionary. National Institute of Health. [www.merriam-webster.com/medlineplus](http://www.merriam-webster.com/medlineplus). March 22, 2010.

Mohamed Jala Khan Interview, 17 year old Secondary School Student at Tudor, April 22<sup>nd</sup>, 2010. SIT Mombasa office.

Mrs. Thirika Interview, April 23<sup>rd</sup>, 2010. Coast Girls Secondary School.

M. Weber, K. Narayan. "Preventing type 2 diabetes: Genes or lifestyle?" *Primary Care Diabetes* 2 (2008): 65-66.

N/A "Kenya: Experts sound alert over rising cases of diabetes among youth." *The Nation*, 30, June, 2007.

N/A "The changing spectrum of Type 2 Diabetes Mellitus." *East African Medical Journal* (August, 2002): 397-399.

Nabila Mohammed Interview, April 20<sup>th</sup>, 2010. Mombasa Hospital.

National Institute of Health. "Diabetes Statistics 2007." April 24, 2010  
<<http://diabetes.niddk.nih.gov/DM/PUBS/statistics/>>

NFL. "NFL Play 60: the NFL Movement for an Active Generation." April 29<sup>th</sup>, 2010. <  
<http://www.nfl.com/news/story?id=09000d5d80b4a489&template=with-video&confirm=true>>



Omran, A. "The epidemiologic transition: a theory of the epidemiology of population change." *Milbank Memorial Fund* 49 (1971): 509-538.

Roglic G, Unwin N, Bennett PH, Mathers C, Tuomilehto J, Nag S, Connolly V, King H. "**The burden of mortality attributable to diabetes: realistic estimates for the year 2000.**" *Diabetes Care* 28 (2005): 2130-2135.

World Health Organization. "Diabetes Key Facts." March 23, 2010  
<<http://www.who.int/mediacentre/factsheets/fs312/en/index.html>>.

Tuomilehto, J. "Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance." *The New England Journal of Medicine* 344 (May, 2001): 1343-1350.

Weight-Control Information Network. "Statistics Related to Overweight and Obesity." NIH Publication Number 04-4158. Updated February 2010. April 20<sup>th</sup>, 2010.  
<<http://www.win.niddk.nih.gov/statistics/index.htm>>

Wikipedia. [www.wikipedia.com](http://www.wikipedia.com) March 22, 2010

World Diabetes Foundation. "Diabetes Care in Nairobi Slums." May 4<sup>th</sup>, 2010 <<http://www.globaldiabeteswalk.net/composite-2574.htm>>

World Diabetes Foundation. "Diabetes Education Programme." May 4<sup>th</sup>, 2010 <<http://www.globaldiabeteswalk.net/composite-452.htm>>

World Diabetes Foundation. Quarterly Update, Winter 2009.  
<<http://www.worlddiabetesfoundation.org/composite-3185.htm>>

World Health Organization. "Diabetes Programme, WHO African Region." April 24, 2010.  
<[http://www.who.int/diabetes/facts/world\\_figures/en/index1.html](http://www.who.int/diabetes/facts/world_figures/en/index1.html)>

World Health Organization. "The World Health Report 2000." April 14<sup>th</sup>, 2010.  
<[http://www.who.int/whr/2000/en/whr00\\_en.pdf](http://www.who.int/whr/2000/en/whr00_en.pdf)>

Writing Committee. "Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies." *The Lancet* 373 (March, 2009): 1083-1096.

Yach D, Stuckler D, Brownell KD. "**Epidemiologic and economic consequences of the global epidemics of obesity and diabetes.**" *Nat Med* 12 (2006): 62-66.

Zimmet, P. "Globalization , coca-colonization and the chronic disease epidemic: can the doomsday scenario be averted?" *Journal of Internal Medicine* 247 (2000): 301-310.